

Evaluating the allelopathic efficiency of the seed powder of *Raphanus sativus* L. in controlling some weeds associating *Phaseolus vulgaris* L.

Ahmed S.A.A., R.R. El-Masry, Nadia K. Messiha and Kowther G. El-Rokiek

Botany Department, National Research Centre, El Buhouth St., Dokki, Giza, Egypt, P.O. Box, 12622.

Received: 11 July 2018 / Accepted: 30 August 2018/ Publication date: 15 Sept. 2018

ABSTRACT

Two pot experiments were conducted in the greenhouse of the National Research Centre, Dokki, Giza, Egypt, in the two successive seasons of 2016 and 2017 to study the allelopathic potentiality of *Raphanus sativus* seed powder (Rssp) on *Phaseolus vulgaris* growth, green and dry yield as well as its effect on the growth of associated weeds i.e. *Corchorus olitorius*, *Abelmoschus esculentus* and *Portulaca oleracea*. Treatments were applied by the incorporation of *R. sativus* seed powder (Rssp) to the soil at (0, 15, 30, 45 and 60 g/kg soil). The results indicated significant reduction in the dry weight of *C. olitorius*, *P. oleracea* and *A. esculentus* in comparison to their corresponding untreated control. The phytotoxic effects of (Rssp) on the three weeds dry weight reached its maximum effect (100%) at 60g / kg soil. The results also indicated that all growth parameters of *P. vulgaris* as well as green and dry yield were significantly increased by different (Rssp) concentrations at the two ages of growth (40 DAS and at harvest) as compared to the corresponding untreated controls. (Rssp) at 30 g/kg soil recorded the highest values as compared to their corresponding controls. The presence of glucosinolates and Phenolic compounds in (Rssp) play an important role in its natural selective bioherbicidal properties in controlling the three weeds associating *P. vulgaris* plants.

Keywords: *Corchorus olitorius*, *Abelmoschus esculentus*, *Portulaca oleracea*, common bean, Allelopathy, *Raphanus sativus*, Glucosinolates content, Phenolic content

Introduction

Phaseolus vulgaris is one of the major grain legumes mostly grown in the world as a source of proteins for human. Like many other grain legumes and crop plants, weeds grown associating these crop plants caused high percentage of yield loss (El-Rokiek *et al.*, 2013 and 2016 and Ahmed *et al.*, 2014). Due to the competition of weeds for nutrients, water, light, etc. different herbicides were used for weed control to decrease its competition and consequently increase the crop yield (Abdelhamid and El-Metwally, 2008 and El-Rokiek *et al.*, 2013).

However, the continuous use of herbicides caused problems because of environmental, toxicological or economical purposes associated with their use (Duke *et al.*, 1999). So, the need for alternative natural herbicides becomes important to reduce the continuous use of synthetic herbicides and for the development of safer, alternative crop protectants (Mahmood and Cheema, 2004). Many crop, trees and weed species have been reported to possess allelopathic activity on the growth of other plant species (Jabran *et al.*, 2010). Allelopathic substances (secondary plant metabolites) are present in many plant tissues, e.g. leaves, stems, flowers, fruits, seeds and roots (Mahmood *et al.*, 2010 and Ahmed *et al.*, 2014). Allelochemicals are released to the environment from plants through degradation, volatilization, leaching from plant leaves, and from root exudation (Petersen *et al.*, 2001 and Price *et al.*, 2005).

The allelopathic potentiality of Brassicaceae plants in suppressing the growth of different types of weeds (annual, perennial and parasitic) have been recently reported from the Botany department of the National Research Centre of Egypt (Messiha *et al.* 2013& 2018; Ahmed *et al.* 2014& 2016; El-Masry *et al.* 2015 and El-Rokiek *et al.* 2017).

Corresponding Author: Ahmed, S.A.A., Botany Department, National Research Centre, El-Buhoth St., Dokki, Giza, Egypt, P.O. Box, 12622.
E-mail: dr_salah_54@yahoo.com

Brassicaceae plant tissues is known to contain considerable amounts of glucosinolates in their tissues which could be easily hydrolyzed in the soil to phytotoxic products such as isothiocyanates, nitriles, thiocyanates, epithionitriles and oxazoliolines (Bones and Rossiter 2006). Isothiocyanates is the main phytotoxic product (Fahey *et al.*, 2001; Zaji and Majd, 2011 and Martinez-Ballesta *et al.*, 2013) and have distinct pesticidal activities (Velasco *et al.*, 2008).

The aim of the present work is to continue our strategy to evaluate the allelopathic efficiency of another member in the Brassicaceae family (*Raphanus sativus*) in controlling weeds associating *Phaseolus vulgaris* plants.

Materials and Methods

Two pot experiments were carried out during two successive seasons of (2016) and (2017) in the greenhouse of the National Research Centre, Dokki, Giza, Egypt. Common bean (*Phaseolus vulgaris*) cv. Giza 4 seeds as well as seeds of Radish (*Raphanus sativus*), Jew's mallow (*Corchorus olitorius*), Okra (*Abelmoschus esculentus*) and Purslane (*Portulaca oleracea*) were obtained from Agriculture Research Centre, Giza, Egypt. Clean seeds of *R. sativus* were grinded to fine powder after that the powder was immediately incorporated in the soil surface before sowing *P. vulgaris* seeds at rate of 0, 15, 30, 45 and 60 g/kg soil. In the same time, the seeds of *P. vulgaris*, *C. olitorius*, *A. esculentus* and *P. oleracea* were sown 2cm deep in plastic pots filled with 2kg of soil. The experiment consisted of 18 treatments including control; each treatment consisted of 8 replicates. All pots were distributed in a complete randomized design. The normal cultural practices of growing *P. vulgaris* plants were followed especially fertilization and irrigation.

Characters studied

Weeds

Three replicates were collected from each treatment at 40 days after sowing (DAS) and at harvest. The dry weight of each weed species was recorded (g /pot) at the two growth ages and the percentage of reduction was calculated as compared to control.

Phaseolus vulgaris plants

Plant growth

Samples of *P. vulgaris* plants at 40 DAS and at harvest were collected from each treatment to determine: plant height (cm), number of leaves/plant and dry weight of plant (g).

Yield and yield components

At harvest, samples of *P. vulgaris* plants were taken from each treatment to determine:
A- Green yield, i.e. number of pods/plant and weight of pods/plant (g)
B- Dry yield, i.e., number of pods/plant, weight of seeds/pod (g), weight of 100 seeds (g) and weight of seeds/plant (g).

Chemical analysis

Total glucosinolates (μ mol/g DW)

Total glucosinolates were extracted from dry samples of seed powder of *R. sativus*. Glucosinolates were measured by determining the liberated glucose which released during hydrolysis by myrosinase enzyme (Rauchberger *et al.*, 1979). The resulting glucose was determined colorimetrically according to the methods defined by Nasirullah and Krishnamurthy (1996).

Total phenolic contents (mg/g DW)

Total phenolic contents of *R. sativus* seeds were determined colorimetrically using Folin and Ciocalteu phenol reagent according to the method defined by Snell and Snell (1953).

Statistical analysis

All data were statistically analyzed according to Snedecor and Cochran (1980) and the treatment means were compared by using LSD at 5% probability.

Results

Weeds growth parameters

The results in Table (1) showed the effect of incorporating different rates from the seed powder of *Raphanus sativus* (0, 15, 30, 45 and 60 g per pot) on the dry weight of the three different weeds i.e. *Corchorus olitorius*, *Abelmoschus esculentus* as well as *Portulaca oleracea* associating the growth of *Phaseolus vulgaris* after 40 DAS and at harvest. The results indicate that the competition between *P. vulgaris* and each weed caused significant decrease in the dry weight of weeds after 40 DAS and lasted till harvest. It is worthy to mention that incorporating the seed powder of *R. sativus* at the rate of 30g/kg soil showed its bioherbicidal efficiency in controlling the different three weeds which reached more than 90% of control. Higher amounts from (Rssp) showed also higher bioherbicidal effect which reached to 100% to the three weeds with the highest rate (60g/kg soil).

Table 1: Effect of incorporating different concentrations of the seed powder of *Raphanus sativus* on dry weight of *Corchorus olitorius*, *Hibiscus esculentus* as well as *Portulaca oleracea* (g/pot).(Combined analysis of the two seasons).

Treatments		At 40 days after sowing		At harvest	
Plants	Concentrations of <i>Raphanus sativus</i> (g/kg soil)	Dry weight of weed (g/pot)	% of reduction	Dry weight of weed (g/pot)	% of reduction
<i>Corchorus olitorius</i>	11.96	00.0	22.85	00.0
<i>Corchorus olitorius</i> + <i>Phaseolus vulgaris</i>	7.45	37.7	19.34	15.4
<i>Corchorus olitorius</i> + <i>Phaseolus vulgaris</i>	15 g	2.41	79.8	3.62	84.2
	30 g	1.63	86.4	1.95	91.5
	45 g	0.82	93.1	1.10	95.2
	60 g	0.43	96.4	0.00	100
<i>Abelmoschus esculentus</i>	13.42	00.0	25.16	00.0
<i>Abelmoschus esculentus</i> + <i>Phaseolus vulgaris</i>	9.69	27.8	21.04	16.4
<i>Hibiscus esculentus</i> + <i>Phaseolus vulgaris</i>	15 g	3.62	73.0	4.15	83.5
	30 g	1.91	85.8	2.27	91.0
	45 g	0.90	93.3	1.41	94.4
	60 g	0.62	95.4	0.00	100
<i>Portulaca oleracea</i>	16.54	00.0	29.40	00.0
<i>Portulaca oleracea</i> + <i>Phaseolus vulgaris</i>	12.37	25.2	24.90	15.3
<i>Portulaca oleracea</i> + <i>Phaseolus vulgaris</i>	15 g	3.95	76.1	8.30	71.8
	30 g	2.06	87.5	6.50	77.9
	45 g	1.08	93.5	1.82	93.8
	60 g	0.86	94.8	0.00	100
LSD at 5%		0.78		1.21	

Phaseolus vulgaris growth

The results in Table (2) indicated the effect of incorporating different amounts of the seed powder of *R. sativus* on different growth parameters of *P. vulgaris* as Plant height (cm), No. of leaves/plant and Dry weight of plant (g) at 40 DAS and at harvest. The results indicated that all growth parameters of *P. vulgaris* were significantly increased by different concentrations of *R. sativus* seed powder (Rssp). Not only, all treatments of Rssp concentrations (15 to 60 g/kg soil) alleviated the harmful effect of the three weeds (*C. olitorius*, *A. esculentus* and *P. oleracea*) associating *P. vulgaris* plants, but also induced significant increase in all *P. vulgaris* growth parameters at the two ages of growth (40 DAS and at harvest) as compared to the corresponding untreated controls. Maximum increase in all *P. vulgaris* growth parameters at 40 DAS and at harvest was recorded with 30g/kg soil Rssp concentration when compared to corresponding controls. The increase in the dry weight of *P. vulgaris* associated with *C. olitorius* or *A. esculentus* or *P. oleracea* at 40 DAS reached to about 10.0, 15.7 and 4.8 %, while at harvest were about 22.6, 31.4 and 16.2 %, respectively over the dry weight of *P. vulgaris* free from weeds as shown from the results in Table (2).

Table 2: Effect of incorporating different concentrations of the seed powder of *Raphanus sativus* on different growth parameters of *Phaseolus vulgaris*. (Combined analysis of the two seasons).

Treatments		Growth parameters					
		At 40 days after sowing			At harvest		
Plants	Concentrations of <i>Raphanus sativus</i> (g/kg soil)	Plant height (cm)	No. of leaves/plant	Dry weight of plant (g)	Plant height (cm)	No. of leaves/plant	Dry weight of plant (g)
<i>Phaseolus vulgaris</i> only	33.2	6.2	2.10	42.5	11.0	4.21
<i>Phaseolus vulgaris</i> + <i>Corchorus olitorius</i>	24.0	3.3	1.10	33.2	5.5	1.79
<i>Phaseolus vulgaris</i> + <i>Corchorus olitorius</i>	15 g	32.0	6.0	1.73	39.2	10.3	3.76
	30 g	35.0	7.8	2.31	49.6	12.6	5.16
	45 g	31.0	5.7	1.61	38.9	10.0	3.57
	60 g	28.0	5.0	1.33	35.8	7.4	2.36
<i>Phaseolus vulgaris</i> + <i>Abelmoschus esculentus</i>	26.0	3.8	1.18	34.0	6.0	1.95
<i>Phaseolus vulgaris</i> + <i>Abelmoschus esculentus</i>	15 g	33.0	6.4	1.84	41.8	10.7	4.13
	30 g	36.0	8.0	2.43	51.5	13.5	5.53
	45 g	33.5	6.7	1.80	44.0	11.5	4.58
	60 g	29.7	4.4	1.50	37.2	9.0	3.29
<i>Phaseolus vulgaris</i> + <i>Portulaca oleracea</i>	22.0	2.7	1.04	32.6	5.2	1.55
<i>Phaseolus vulgaris</i> + <i>Portulaca oleracea</i>	15 g	30.0	5.5	1.56	38.0	9.6	3.40
	30 g	34.0	7.3	2.20	47.3	12.0	4.89
	45 g	29.0	4.6	1.45	36.4	8.0	3.07
	60 g	26.5	4.1	1.24	35.0	7.0	2.15
LSD at 5%		1.91	0.88	0.81	1.93	1.18	1.08

Phaseolus vulgaris yield

A- Green yield

The results of the green yield of *P. vulgaris* associated with *C. olitorius* or *A. esculentus* or *P. oleracea* recorded in Table (3) cleared that different concentrations of Rssp (15 to 60 g/kg soil) induced significant increase in the number of green pods/plant and weight of pods/plant of *P. vulgaris* when compared to their corresponding controls. Maximum increase in number and weight of green

Pods/plant of *P. vulgaris* was recorded with 30 g/kg soil Rssp treatment as compared to their corresponding controls. Not only this treatment (Rssp at 30 g/kg soil) alleviated the reduction caused by the effect of *C. olitorius* or *A. esculentus* or *P. oleracea* on the weight of green pods/plant that reached to about 72.2, 68.3 and 74.8%, respectively, but also increased this character to about 7.6, 9.9 and 3.7%, respectively over their corresponding healthy control.

Table 3: Effect of incorporating different concentrations of the seed powder of *Raphanus sativus* on yield and yield components of *Phaseolus vulgaris*. (Combined analysis of the two seasons).

Treatments		Yield and yield components					
Plants	Concentrations of <i>Raphanus sativus</i> (g/kg soil)	Green yield		Dry yield			
		No. of green pods/plant	Weight of green pods/plant (g)	No. of dry pods/plant	Weight of seeds/pod (g)	Weight of 100 seeds (g)	Weight of seeds /plant (g)
<i>Phaseolus vulgaris</i> only	7.20	13.74	7.36	1.64	29.7	10.12
<i>Phaseolus vulgaris</i> + <i>Corchorus olitorius</i>	3.30	3.82	3.16	0.86	21.0	3.45
<i>Phaseolus vulgaris</i> + <i>Corchorus olitorius</i>	15 g	6.30	10.10	6.14	1.49	28.9	9.51
	30 g	8.32	14.79	8.01	1.89	33.2	12.74
	45 g	6.00	10.30	5.95	1.38	28.0	9.36
	60 g	4.36	6.50	3.99	1.10	23.8	5.25
<i>Phaseolus vulgaris</i> + <i>Abelmoschus esculentus</i>	3.52	4.36	3.23	0.95	21.8	3.97
<i>Phaseolus vulgaris</i> + <i>Abelmoschus esculentus</i>	15 g	6.90	11.25	6.61	1.56	31.1	10.60
	30 g	8.65	15.10	8.20	1.93	34.3	13.10
	45 g	7.60	12.32	6.95	1.70	31.6	11.42
	60 g	5.51	7.42	4.97	1.22	25.3	7.83
<i>Phaseolus vulgaris</i> + <i>Portulaca oleracea</i>	2.95	3.46	2.96	0.79	20.0	2.91
<i>Phaseolus vulgaris</i> + <i>Portulaca oleracea</i>	15 g	5.63	8.80	5.72	1.26	27.0	8.74
	30 g	8.00	14.25	7.84	1.77	32.7	12.10
	45 g	5.10	8.40	4.19	1.19	24.3	7.53
	60 g	4.20	5.80	3.62	1.00	23.0	4.06
LSD at 5%		1.03	1.16	1.15	0.65	2.08	1.35

B- Dry yield

Dry yield and yield components of *P. vulgaris* associated with *C. olitorius* or *A. esculentus* or *P. oleracea* cleared that all Rssp concentrations (15 to 60 g/kg soil) induced significant increase in the different yield parameters (number of dry pods/plant, weight of seeds/plod, weight of 100 seeds as well as weight of seeds/plant) as compared to corresponding untreated controls (Table 3). The best treatment was recorded with 30g/kg soil Rssp concentration that achieved the highest increase in all *P. vulgaris* plant yield parameters. The increase in weight of seeds/plant of *P. vulgaris* associated with *C. olitorius* or *A. esculentus* or *P. oleracea* reached to about 25.9, 29.4 and 19.6 %, respectively over their corresponding control free from weeds.

Table 4: Total glucosinolates ($\mu\text{mol/g}$ dry weight) and Total phenolic contents (mg/g dry weight) in the seed powder of *Raphanus sativus*

Material	Total glucosinolates($\mu\text{mol/g}$ dry weight)	Total phenolic contents(mg/g dry weight)
Seed powder of <i>Raphanus sativus</i>	688.54	69.50

Discussion

The allelopathic compounds (allelochemicals) released from plants into the environment, as a result of secondary metabolites, include a variety of compounds, often attract or repel, nourish or poison to other organisms. Allelochemicals like phenolic compounds, flavonoids, terpenoids, alkaloids, amino acids and glucosinolates were found in allelopathic plants (Fahey *et al.*, 2001; Velasco *et al.*, 2008 and Ahmed *et al.*, 2012).

Recently, several researches showed the potentiality of using the allelopathic technique as a component of integrated weed management as bioherbicide to suppress weeds in crops (Zaji and Majd, 2011; Ahmed *et al.*, 2012, 2016; Messiha *et al.*, 2013 & 2018; El-Masry *et al.*, 2015 and El-Rokiek *et al.*, 2017). Also, weed management systems seek biological solutions to minimize the harmful effects resulted from the use of herbicides in agricultural systems. Therefore, allelochemicals could be considered as an important tool for sustainable weed control management (El-Metwally *et al.*, 2014 and El-Wakeel, 2015).

The results of the present investigation reveal to great extent significant reduction in the three weeds i.e. *Corchorus olitorius*, *Abelmoschus esculentus* and *Portulaca oleracea* growth after the incorporation of (Rssp) to the soil till 40 (DAS). Complete reduction of all weeds recorded by the higher concentration (60g/kg soil) of (Rssp) at harvest (Table 1). The previous results showed that Brassicaceae family has allelopathic potential on the growth of other plants (Petersen *et al.*, 2001; Messiha *et al.*, 2013&2018; Ahmed *et al.*, 2014& 2016; Bashen, 2014 and El-Masry *et al.*, 2015). In this connection, it is worthy to mention that the allelopathic effects of Brassicaceae plants were attributed to its natural allelochemicals mainly glucosinolates and phenolic compounds (Table 4). Glucosinolates hydrolyzed by endogenous enzyme myrosinase to a number of products. The main breakdown products are isothiocyanates, which are phytotoxic and achieved good results in controlling weeds (Zaji and Majd, 2011; Martinez-Ballesta *et al.*, 2013; Messiha *et al.*, 2013& 2018 ; Ahmed *et al.*, 2014,&2016; El-Masry *et al.*, 2015 and El-Rokiek *et al.*, 2017). Moreover, Petersen *et al.*, 2001 and Uremis *et al.*, 2009 reported that the allelopathic effect of *Raphanus sativus* L. could be also due to the presence of p-hydroxy benzoic acid in addition to isothiocyanates.

On the other side (Rssp) treatments not only achieved to great extent good results in controlling the three weeds i.e. *C. olitorius* (C₃ plant), *A. esculentus* (C₃ plant) and *P. oleracea* (C₄ plant) but also increased *P. vulgaris* growth (Table 2) and consequently improved its green and dry yield (Table 3). Several workers found that the inhibition of weed growth increased the competitive ability of the crop plant and consequently improved growth and yield (Abdelhamid and El-Metwally, 2008; Ahmed *et al.*, 2012 & 2014; El-Rokiek *et al.*, 2013 and El-Masry *et al.*, 2015).

It is worthy to mention that (Rssp) at 30g/kg soil was the best treatment in controlling the three weeds (*C. olitorius*, *A. esculentus* and *P. oleracea*) and consequently increased the growth and yield of *P. vulgaris* as compared to corresponding controls, this may be due to the selectivity of allelochemicals in their action and plants in their responses (Einhellig, 1995). Allelochemicals which inhibit the growth of some species at certain concentrations may stimulate the growth of same or different species at different concentrations (Ahmed *et al.*, 2012 & 2014; Messiha *et al.*, 2013& 2018 and Bashen, 2014).

The results of the present work indicate clearly the possibility of using allelopathic activity of *Raphanus sativus* seed powder as a selective bioherbicide for controlling annual weeds.

References

- Abdelhamid, M.T. and I.M. El-Metwally, 2008. Growth, nodulation and yield of soybean and associated weeds as affected by weed management. *Planta Daninha* 26 (4): 855-863.
- Ahmed, S.A., K.G. El-Rokiek, R.R. El-Masry and N.K. Messiha, 2014. The Efficiency of Allelochemicals in The Seed Powder of *Eruca sativa* in Controlling Weeds in *Pisum sativum*. *Middle East Journal of Agriculture Research*, 3(4): 757-762.
- Ahmed, S.A.A., N.K. Messiha, K.G. El-Rokiek, S.A. Mohamed and R.R. El-Masry, 2016. The allelopathic efficiency of two Brassicaceae plant seeds in controlling weeds associating sunflower plants. *Res. J. of Pharm., Biol. and Chem. Sci.* Sept.-Oct., 7(5): 158-165.

- Ahmed, S.A., N.K. Messiha, R.R. El-Masry and K.G. El-Rokiek, 2012. Allelopathic potentiality of the leaf powder of *Morus alba* and *Vitis vinifera* on the growth and propagative capacity of purple nutsedge (*Cyperus rotundus* L.) and maize (*Zea mays* L.). J. of Appl. Sci. Res., 8(8): 4744-4751.
- Bashen, A.A., 2014. Morphological and elements constituent effects of Allelopathic Activity of some medicinal plants extracts on *Zea mays*. Int. J. Curr. Res. Aca. Rev., 2(4):135-145.
- Bones, A.M. and J.T. Rossiter, 2006. The enzymic and chemically induced decomposition of glucosinolates. Phytochemistry, 67: 1053-1067.
- Duke, S. O., F.E. Fedayan, J.G. Romagni and A.M. Rimando, 1999. Natural Products as sources of herbicides: current status and future trends. Weed Res., 40: 99-111.
- Einhellig F.A., 1995. Mechanism of Action of Allelochemical in Allelopathy. In Allelopathy Organisms, Processes and Application. Am. Chem. Soc., Washington, USA, pp.: 96-116.
- El-Masry, R.R., N.K. Messiha, K.G. El-Rokiek, S.A. Ahmed and S.A. Mohamed, 2015. The Allelopathic Effect of *Eruca sativa* Mill. Seed Powder on Growth and Yield of *Phaseolus vulgaris* and Associated Weeds. Current Sciences International, 4(4):485-490.
- El-Metwally, I.M., M.N. Shehata, E.R. El-Desoki, S.A. Ahmed and M.A. El-Wakeel, 2014. Response of Wheat and Associated Weeds to Allelopathic Water Extracts Tank Mixed with Reduced Wheat Herbicides Dose. Middle East J Appl. Sci., 4(4): 873-883.
- El-Rokiek, K.G., E.R. El-Desoki, I.M. El-Metwally and M.G. Dawood, 2016. Allelopathic Effect of Mango Leaf Residue against *Portulaca oleracea* or *Corchorus olitorius* Associated *Phaseolus vulgaris* Growth. Research Journal of Pharmaceutical, Biological and Chemical Sciences, 7(4): 964-970.
- El-Rokiek, K.G., M.T. Abdelhamid and S.A. Saad El-Din, 2013. Physiological response of purslane weed (*Portulaca oleracea*) and two common beans (*Phaseolus vulgaris*) recombinant inbred lines to Phosphorus fertilizer and bentazon herbicide. Journal of Applied Sciences Research, 9(4): 2743-2749.
- El-Rokiek, K.G., S.A.A. Ahmed, N.K. Messiha, S.A. Mohamed and R.R. El-Masry, 2017. Controlling the Grassy weed *Avena fatua* associating wheat plants with the seed powder of two brassicaceae plants *Brassica rapa* and *Sinapis alba*. Middle East J Agri. Res., 6(4):1014-1020.
- El-Wakeel, M.A.M., 2015. Effect of Allelopathy and Autotoxicity of Some Plants As Well As Herbicides on Wheat and Associated Weeds. Ph.D. Thesis, Fac. Of Sci., Benha Univ., Cairo.
- Fahey, J.W., A.T. Zalczman and P. Talalay, 2001. The chemical diversity and distribution of glucosinolates and isothiocyanates plants. Phytochemistry, 56: 5-51.
- Jabran K., M. Farooq, M. Hussain, R. Hafeez-ur and M.A. Ali, 2010. Wild oat (*Avena Fatua* L.) and Canary Grass (*Phalaris Minor* Ritz.) Management through Allelopathy. J. Plant Protec. Res., 50 (1): 41-44.
- Mahmood, A. and Z.A. Cheema, 2004. Influence of sorghum mulch on purple nutsedge (*Cyperusrotundus* L.). Int. J. Agric. & Biol., 6(1): 86-88.
- Mahmood, A., Z.A. Cheema, A. Khaliq and A.ul-Hassan, 2010. Evaluating the potential of allelopathic plant water extracts in suppressing horse purslane growth. Int. J. Agric. & Biol., 12 (4): 581-585.
- Martinez-Ballesta, M., D.A. Moreno and M. Carvajal, 2013. The Physiological Importance of Glucosinolates on Plant Response to Abiotic Stress in Brassica. Int. J. Mol. Sci., 14: 11607-11625.
- Messiha, N.K., M.A.T. El-Dabaa, R.R. El-Masry and S.A.A. Ahmed, 2018. The allelopathic influence of *Sinapis alba* seed powder (white mustard) on the growth and yield of *Vicia faba* (faba bean) infected with *Orobanche crenata* (broomrape). Middle East Journal of Applied Sciences, 8(2): 418-425.
- Messiha, N.K., S.A. Ahmed, K.G. El-Rokiek, M.G. Dawood and R.R. El-Masry, 2013. The Physiological Influence of Allelochemicals in Two Plant Seeds on the Growth and Propagative Capacity of *Cyperus rotundus* L. and *Zea mays* L. World Applied Sciences Journal, 26(9):1142 – 1149.
- Nasirullah and M.N. Krishnamurthy, 1996. A method for estimating glucosinolates in mustard/rape seeds and cake. J. Sci. Technol., 33(6):498-500.
- Petersen, J., R. Betz, F. Walker and K. Hurler, 2001. Weed suppression by release of isothiocyanates from turnip-rape mulch. Agronomy Journal (93):37-42.

- Price, A.J., C.S. Charron and C.E. Sams, 2005. Allyl isothiocyanate and carbon dioxide produced during degradation of *Brassica juncea* tissue in different soil conditions. Hort. Sci., 40:1734-1739.
- Rauchberger, Y., S. Mokady and U. Cogan, 1979. The effect of aqueous leaching of glucosinolates on the nutritive quality of rapeseed meal. J. Food Agric., 30: 31-39.
- Snedecor, G.W. and W.G. Cochran, 1980. Statistical Methods. 7th Ed. pp.: 507. The Iowa State Uni. PRESS, Ames, Iowa.
- Snell, F.D. and C.T. Snell, 1953. Colorimetric Methods. Pp.:66 Volume 111. Organi, D. Van Nostrand Company, In R.E.C. Toronto, New York, London.
- Uremis, I., M. Arslan, A. Uludag and M. Sangun, 2009. Allelopathic potentials of residues of 6 Brassica species on Johnson grass (*Sorghum halepense* (L.) Pers.). African Journal of Biotechnology, 8: 3497-3501.
- Velasco, P., P. Soengas, M. Vilar and M.E. Cartea, 2008. Composition of glucosinolates profiles in leaf and seed tissue of different *Brassic napus* crop. J. Amer. Soc. Hort. Sci., 133 (4): 551-558.
- Zaji, B. and A. Majd, 2011. Allelopathic potential of canola (*Brassica napus* L.) residues on weed suppression and yield response of maize (*Zea mays* L.). International Conference on Chemical, Ecology and Environmental Sciences IICCEES, 2011) Pattaya, December: 457-460.